

EMERGENCY MEDICINE ALERT[®]

An essential monthly update of developments in emergency medicine

From the Publishers of Emergency Medicine Reports[™]

Enclosed in this issue:
Trauma Reports

Thomson American Health Consultants Home Page—<http://www.ahcpub.com>

CME for Physicians—<http://www.cmeweb.com>

THOMSON
AMERICAN HEALTH
CONSULTANTS

INSIDE

*Subcutaneous
lispro insulin
for diabetic
ketoacidosis*
page 50

*ED
thoracotomy
indications
in the trauma
patient*
page 51

*ACE-inhibitor
angioedema:
How common?
How serious?*
page 52

*Special Feature:
Heparin use
in acute
coronary
syndromes*
page 53

Bedside Ultrasound to Confirm Pediatric Endotracheal Tube Placement

ABSTRACT & COMMENTARY

Source: Hsieh KS, et al. Secondary confirmation of endotracheal tube position by ultrasound image. *Crit Care Med* 2004; 32:S374-377.

VERIFICATION OF ENDOTRACHEAL TUBE PLACEMENT IN THE AIRWAY following intubation is critical in the emergent setting. In addition to auscultation and other clinical assessments, new techniques for tube confirmation include measurement of end-tidal carbon dioxide (CO₂) and use of the esophageal detector device. However, each of these methods has limitations, and thus, multiple methods of real-time confirmation are recommended.

Investigators studied the utility of ultrasonography to verify correct placement of the endotracheal tube in patients cared for in a pediatric intensive care unit (ICU). During a six-month study period, they studied 59 patients who were intubated for either cardiopulmonary arrest or impending respiratory failure. Ultrasound imaging was performed using a portable machine with 2-D and M-mode scanning and transducer frequencies varying from 2.5 to 7.5 MHz. Several scanning windows were utilized, primarily the subxiphoid window with a wide scanning angle (70-85°) to allow visualization of both sides of the diaphragm. Operators assessed diaphragmatic movement to confirm tube placement. If both sides of the diaphragm moved toward the abdomen during positive ventilation and toward the chest during exhalation, then the tube was determined to be placed correctly. Opposite movement indicated esophageal placement. Asymmetric movement of the left and right sides of the diaphragm indicated tube placement in a mainstem bronchi.

The authors report that this ultrasound technique correctly identified two esophageal intubations as well as eight right mainstem bronchus intubations. Because confirmation was performed real-time at the bedside, the investigators reported that these misplacements were corrected rapidly without difficulty. Moreover, for the right mainstem intubations, the endotracheal tubes were withdrawn a small distance until bilateral equal motion of the diaphragm could be seen clearly on ultrasound. Subsequently, the authors report that their

EDITOR

Richard A. Harrigan, MD, FAAEM
Associate Professor of Emergency Medicine, Temple University Hospital and School of Medicine, Philadelphia, PA

EDITORIAL BOARD

Stephanie B. Abbuhl, MD, FACEP
Medical Director, Department of Emergency Medicine, The Hospital of the University of Pennsylvania; Associate Professor of Emergency Medicine, University of Pennsylvania School of Medicine, Philadelphia

William J. Brady, MD

Associate Professor of Emergency Medicine and Internal Medicine, Vice Chair, Emergency Medicine, University of Virginia, Charlottesville

Theodore C. Chan, MD, FACEP

Associate Clinical Professor of Medicine, Emergency Medicine, University of California, San Diego

Michael Felz, MD

Associate Professor, Department of Family Medicine, Medical College of Georgia, Augusta

Ken Grauer, MD

Professor and Associate Director, Family Practice Residency Program, Department of Community Health and Family Medicine, College of Medicine, University of Florida, Gainesville

Richard J. Hamilton, MD, FAAEM, ABMT

Associate Professor of Emergency Medicine, Residency Program Director, Department of Emergency Medicine, Drexel University College of Medicine, Philadelphia, PA

David J. Karras, MD, FAAEM, FACEP

Associate Professor of Emergency Medicine, Associate Chair for Academic Affairs, and Research Director, Department of Emergency Medicine, Temple University School of Medicine, Philadelphia, PA

Andrew D. Perron, MD, FACEP, FACS

Residency Program Director, Department of Emergency Medicine, Maine Medical Center, Portland, ME

Jacob W. Ufberg, MD

Assistant Professor of Emergency Medicine, Residency Program Director, Department of Emergency Medicine, Temple University School of Medicine, Philadelphia, PA

Special Clinical Projects and Medical Education Resources:

Gideon Bosker, MD

Assistant Clinical Professor, Section of Emergency Services, Yale University School of Medicine, Associate Clinical Professor, Oregon Health Sciences University, Portland, OR

ultrasound technique confirmed all 59 intubations. They conclude that ultrasound imaging of diaphragm motion allows a useful, quick, noninvasive, real-time anatomic method for assessment and confirmation of endotracheal tube position. ❖

■ COMMENTARY BY TED CHAN, MD, FACEP

Bedside ultrasonography performed by the ED physician is becoming more common for many indications. This study is an interesting report on using ultrasound as another means of verifying endotracheal tube placement rapidly at the bedside. Whereas others have attempted to actually visualize the upper airway during intubation with limited success,¹ this study uses ultrasound to visualize the diaphragm to assess airway control.

The authors report excellent success with their approach, even suggesting that real-time ultrasound can

provide gross tuning for correcting mainstem intubations. However, a number of limitations should be kept in mind when considering this study. First, the total number of patients was small, and a much larger study is needed to confirm ultrasonography's applicability in this setting. Second, the gold standard to determine correct tube location is not clear in this study, which could have affected the results. Third, the study population was limited to patients in a pediatric ICU; it is not clear how well this technique would perform in adults, particularly those with greater soft tissue, which may interfere with sonographic imaging. Finally, as this technique is based on diaphragmatic movement, it is not clear if there could be differences in assessing motion in patients who are paralyzed compared with those who are spontaneously breathing when intubated.

This concept is intriguing, but more study is needed. Although it is unlikely that ultrasonography will become the single gold-standard method for verifying tube placement any time soon, it could add to our current armamentarium of methods for intubation confirmation.

Reference

1. Ma G, et al. Using ultrasound to visualize and confirm endotracheal intubation. *Acad Emerg Med* 1999;6:515.

Emergency Medicine Alert, ISSN 1075-6914, is published monthly by Thomson American Health Consultants, 3525 Piedmont Rd., NE, Bldg. 6, Suite 400, Atlanta, GA 30305.

Vice President and Group Publisher: Brenda Mooney.
Editorial Group Head: Valerie Loner.
Managing Editor: Martha Jo Dendinger.
Marketing Manager: Schandale Kornegay.

GST Registration Number: R128870672.

Periodicals postage paid at Atlanta GA 30304. **POSTMASTER:** Send address changes to **Emergency Medicine Alert**, P.O. Box 740059, Atlanta, GA 30374.

Copyright © 2004 by Thomson American Health Consultants. All rights reserved. No part of this newsletter may be reproduced in any form or incorporated into any information retrieval system without the written permission of the copyright owner.

Back issues: \$48. One to nine additional copies, \$234 each; 10 to 20 additional copies, \$173 each.

This is an educational publication designed to present scientific information and opinion to health professionals to stimulate thought and further investigation. It does not provide advice regarding medical diagnosis or treatment for any individual case. It is not intended for use by the layman.

THOMSON
★
AMERICAN HEALTH CONSULTANTS

Conflict of Interest Disclosure

To reveal any potential bias in this publication, and in accordance with Accreditation Council for Continuing Medical Education guidelines, Drs. Harrigan (editor), Abbuhl, Chan, Felz, Hamilton, Perron, and Ufberg have reported no relationships with companies having ties to the field of study covered by this CME program. Dr. Grauer is sole proprietor of KG/EKG Press. Dr. Karras has reported that he is a consultant for Bayer Pharmaceuticals; consultant, speaker and researcher for Aventis Pharma; and a researcher for Bristol-Myers Squibb and Sepracor Inc. Dr. Brady is on the speaker's bureau for Genentech. Dr. Pierson is on the speaker's bureau of GlaxoSmithKline, Boehringer-Ingelheim, 3-M, Bayer, and Astra Zeneca. Dr. Bosker also acknowledges that he receives royalties, commissions, and other compensation relating to the sale of textbooks, reprints of articles, and other written materials to the following pharmaceutical companies: Pfizer, Genentech, Aventis, Pharmacia, and Bayer. This publication does not receive commercial support.

Subscriber Information

Customer Service: 1-800-688-2421

Customer Service E-Mail Address:

customerservice@thomson.com

Editorial E-Mail Address: martha.dendinger@thomson.com

World-Wide Web: <http://www.ahcpub.com>

Subscription Prices

United States: \$299 per year (Resident rate: \$144.50)

Canada: \$329 per year plus GST (Resident rate: \$159.50)

Elsewhere: \$329 per year (Resident rate: \$159.50)

Accreditation

Thomson American Health Consultants (AHC) is accredited by the Accreditation Council for Continuing Medical Education (ACCME) to provide continuing medical education for physicians. This CME activity was planned and produced in accordance with ACCME Essentials.

Thomson American Health Consultants designates this continuing medical education activity for up to 20 hours in category 1 credit toward the AMA Physician's Recognition Award. Each physician should only claim those hours of credit that he/she actually spent in the educational activity.

Emergency Medicine Alert also is approved by the American College of Emergency Physicians for 20 hours of ACEP category 1 credit.

Emergency Medicine Alert has been approved by the American Academy of Family Physicians as having educational content acceptable for Prescribed credit hours. This volume has been approved for up to 20 Prescribed credit hours. Term of approval covers issues published within one year from the beginning distribution date of June 2003. Credit may be claimed for one year from the date of this issue. **For CME credit, add \$50.**

This CME activity is intended for emergency physicians. It is in effect for 36 months from the date of the publication.

Questions & Comments

Please call **Martha Jo Dendinger**, Managing Editor, (404) 262-5514, or martha.dendinger@thomson.com.

Subcutaneous Lispro Insulin for Diabetic Ketoacidosis

ABSTRACT & COMMENTARY

Source: Umpierrez GE, et al. Efficacy of subcutaneous insulin lispro vs. continuous intravenous regular insulin for the treatment of patients with diabetic ketoacidosis. *Am J Med* 2004; 117: 291-296.

THE STANDARD OF CARE FOR THE TREATMENT OF DIABETIC ketoacidosis (DKA) is hospitalization for administration of regular insulin by continuous intravenous (IV) infusion. Patient monitoring is time- and labor-intensive, and inadvertent hypoglycemia is a recognized complication. In the majority of hospitals, therefore, patients being treated for DKA are admitted to the intensive care unit (ICU) until their hyperglycemia and ketonemia have resolved.

Insulin lispro is an analog of human insulin with a more rapid onset than regular insulin (10-20 min vs 1-2 hr). Insulin lispro also has a faster time to peak activity than regular insulin (about 1 hr vs about 3 hr) and shorter duration of activity. The authors of this study proposed that patients with DKA could be treated safely

in a non-ICU setting using repeated subcutaneous (SQ) doses of insulin lispro, and performed a study comparing this treatment regimen with standard regular insulin therapy administered by continuous IV infusion within the ICU setting.

Patients qualified for the study if they presented to the ED with DKA and were not critically ill. They were randomized to receive therapy with either insulin lispro SQ (lispro group) or regular insulin by IV (regular group). Those in the lispro group were admitted to a general medicine ward or intermediate care unit and received a 0.3 unit/kg SQ injection, followed by an hourly 0.1 U/kg injection. Patients in the regular insulin group received an initial bolus of 0.1 U/kg, followed by a continuous infusion at 0.1 U/kg/hr. Blood glucose levels were checked hourly in the lispro group and every 2 hours in the regular group; serum chemistry was checked every 2 hours initially, then every 4 hours. The endpoint was resolution of hyperglycemia and ketoacidosis.

Twenty patients were enrolled in each study group. There were no intergroup differences in time to correction of hyperglycemia (7 hours in each group) or time to correction of ketoacidosis (10-11 hours). There were also no differences in rates of hypoglycemic events (one patient in each group) or length of hospital stay. Hospital charges were substantially greater for patients in the regular insulin group, who were, by design, admitted to the ICU. The authors conclude that DKA can be safely and economically treated in non-ICU settings using insulin lispro SQ rather than regular insulin IV. ❖

■ COMMENTARY BY DAVID J. KARRAS, MD, FAAEM, FACEP

The value of this study lies in demonstrating the feasibility of treating patients with DKA with SQ injections of fast-acting insulin analogs. By virtue of their rapid onset and shorter half-lives, repeated SQ doses may be used to titrate the blood glucose levels with a low risk of bottoming-out the blood sugar. Thus, these analogs may be appropriate for frequent administration in a non-ICU setting. The authors demonstrate that therapy with SQ insulin lispro resolves DKA as quickly and safely as therapy with IV regular insulin.

The cost savings demonstrated by the study is actually an artifact of the study design. By protocol, ICU admission was mandated for the regular insulin group, while the lispro group was admitted to a regular ward or intermediate care unit. The lower cost of treatment associated with lispro therapy resulted from the higher cost of the ICU stay itself and the comparison is not valid. Also by design, the lispro group actually received more frequent blood glucose checks than the regular insulin group. This might suggest a higher cost of therapy asso-

ciated with lispro had the patients been treated in comparable settings, but again, this is an artifact of study design. In short, the authors cannot render conclusions regarding cost-effectiveness of insulin lispro therapy.

A more persuasive conclusion, entirely justified by the study, is that stable patients with DKA routinely do not require ICU admission, even if they receive regular insulin by continuous IV infusion. I suspect that a shift in the treatment setting would result in far greater cost savings than a change in the type of insulin or method of delivery.

ED Thoracotomy Indications in the Trauma Patient

ABSTRACT & COMMENTARY

Source: Powell DW, et al. Is emergency department resuscitative thoracotomy futile care for the critically injured patient requiring prehospital cardiopulmonary resuscitation? *J Am Coll Surg* 2004;199:211-215.

THE ROLE OF RESUSCITATIVE EMERGENCY DEPARTMENT thoracotomy (EDT) remains controversial; most studies of this procedure have been retrospective with few survivors. To reduce futile care and minimize health care provider risk in resuscitative EDT for patients with prehospital arrest after trauma, the National Association of Emergency Medical Services Physicians Standards and Clinical Practice Committee and the American College of Surgeons Committee on Trauma released guidelines that address the topic. The guidelines state that EDT “does not appear to have a role in prehospital traumatic cardiopulmonary arrest as a result of blunt trauma...” and that traumatic cardiopulmonary arrest “secondary to penetrating trauma, while still having a dismal prognosis, may be more amenable [to EDT] in the case of isolated penetrating trauma to the thorax.”¹

This study questioned whether selected patients who fail to respond to prehospital cardiopulmonary resuscitation (CPR) are potentially salvageable, and should be transported to the nearest trauma center for evaluation and prompt EDT. The study authors prospectively collected a database of all patients receiving EDT during a 26-year study period. The data included mechanism of injury, prehospital and ED vital signs, need for and duration of prehospital CPR, presence or absence of prehospital endotracheal intubation, cardiac rate and rhythm at the time of EDT, survival to hospital discharge, and neurologic outcome deficits (i.e., none, mild, or severe). During the study period, 959 patients underwent EDT with 62 (6.5%) survivors. The injury mecha-

nism for survivors was stab wound in 39 (63%), gunshot wound in 12 (19%), and blunt trauma in 11 (18%). Thirty-three of these patients had no detectable vital signs on arrival to the ED. Twenty-six (42%) of the survivors required prehospital CPR, and were the focus of data analysis.

Among the 26 patients surviving EDT after prehospital CPR, the mean age was 30 ± 3 years and the mean base deficit was 21.5 (range 14 to 25). The mechanism of injury was stab wound in 18 (69.2%), gunshot wound in four (15.4%), and blunt trauma in four (15.4%). Twenty-one of the 26 patients left the hospital with no or mild neurologic deficits (15 none, six mild). Thirteen of the 15 (87%) patients discharged with no neurologic impairment were victims of stab wounds. Of the five patients with severe neurologic deficits, four were associated with a blunt mechanism of injury. Thus, all patients with a blunt mechanism of injury who survived prehospital CPR and EDT had severe neurologic impairment.

Five survivors had asystole documented at the time of EDT. Four of these patients survived with no or mild neurologic deficits, and all had pericardial tamponade due to stab wounds to the ventricle. The initial base deficit in the ED did not appear to predict neurologic outcome. All survivors had successful endotracheal intubation in the prehospital setting and fewer than 15 minutes of prehospital CPR.

The authors submit the following guidelines based on their study. Patients with blunt mechanism of injury should be declared dead on arrival if they have received more than 5 minutes of prehospital CPR and have no signs of life (e.g., pupillary response, motor activity, or respiratory effort). A patient with penetrating mechanism of injury should be declared dead on arrival if he has received more than 15 minutes of prehospital CPR with no signs of life in the ED, or if he has asystole on arrival and does not have a penetrating wound that could result in pericardial tamponade. ❖

■ COMMENTARY BY JACOB UFBERG, MD

A great deal of effort has been devoted to the discussion of when a trauma patient should be declared dead on arrival, and when heroic efforts, such as EDT, are warranted and may yield neurologically functional survival. Unfortunately, we are unlikely to see a large, randomized trial of the use of EDT in the near future. The number of patients who receive EDT in a given center is small, and the number of survivors is far smaller (as evidenced by only 62 survivors in one major trauma center during a 26-year period). Thus, we are forced to rely upon studies such as this one to help us decide when EDT is indicated.

This report echoes previous reports in that the highest EDT survival rates are seen among patients with ventricular injury and pericardial tamponade due to stab wounds. A small number of patients survived other penetrating mechanisms after requiring prehospital CPR. Thus, their recommendations regarding EDT for penetrating trauma seem reasonable.

Unfortunately, the authors do not provide the mechanism of injury for the 897 non-survivors of EDT so that we can draw conclusions on the overall survival rates according to the mechanism of injury. Even without this information, we can see the incredibly small numbers of EDT survivors among patients with blunt mechanism of injury requiring prehospital CPR. We also can see that all of these survivors had dismal neurologic outcomes. The authors base their recommendation of fewer than 5 minutes of prehospital CPR in blunt trauma on sporadic reports of patients with good outcomes who lost vital signs just before arrival. Even with these case reports, it would seem reasonable to withhold EDT in all patients with blunt mechanism of injury and no signs of life on arrival to the ED.

Reference

1. Hopson LR, et al. Guidelines for withholding or termination of resuscitation in prehospital traumatic cardiopulmonary arrest. Joint Position Statement of the National Association of EMS Physicians and the American College of Surgeons Committee on Trauma. *J Am Coll Surg* 2003;196:106-112.

ACE-Inhibitor Angioedema: How Common? How Serious?

ABSTRACT & COMMENTARY

Source: Sondhi D, et al. Airway compromise due to angiotensin-converting enzyme inhibitor-induced angioedema: Clinical experience at a large community teaching hospital. *Chest*. 2004;126:400-404.

IN THIS RETROSPECTIVE STUDY FROM A COMMUNITY teaching hospital in Philadelphia, Sondhi and colleagues identified all patients whose admissions had been assigned the diagnosis code for angioedema (AE) during the five academic years starting in 1996. They reviewed these patients' medical records to determine the presenting symptoms and signs, treatments and outcomes, as well as whether the patients were taking angiotensin-converting enzyme inhibitors (ACEIs) when they developed symptoms. All patients on ACEIs were assumed to have ACEI-induced AE. During the

Table. Findings and Interventions Among 45 Patients Presenting with ACE-Inhibitor-Related Angioedema

Lip and tongue swelling	45 (100%)
Pulmonary manifestations	17 (38%)
Dyspnea	17/17 (100%)
Stridor/respiratory failure	5/17 (29%)
Cough	2/17 (12%)
Dysphagia	9 (20%)
Droping	8 (18%)
Pruritis	6 (13%)
Admitted to ICU	18 (40%)
Intubated	5 (11%)
ICU pts intubated	5/18 (28%)

five-year study period, 70 patients were admitted with AE, among whom this was considered related to ACEI therapy in 45 (64%). In the other 25 patients, AE was ascribed to food allergy, antibiotics, or infection in 23. No precipitating factor could be identified in two cases.

Of the 45 patients with ACEI-related AE, the mean age was 62 years and 41 (91%) were African-American. Twenty-nine (67%) were women. No specific ACEI appeared more likely to cause AE than the others. Duration of ACEI use prior to presentation ranged from 1 day to 5 years; 21 of 45 (45%) patients presented within two months of starting the drug, while 11 (24 patients) had been on it for six months to five years.

Presenting symptoms and signs are shown in the Table. All patients were treated in the ED with H1 blockers, methylprednisolone, and intravenous fluids, and 15 of them received epinephrine. Eighteen patients (40%) were admitted to the ICU, and five (28% of those requiring intensive care) required intubation, one of whom underwent emergency cricothyrotomy. Mean duration of intubation in these patients was 2.2 days, and patients admitted to the ICU spent an average of 2.2 days there. Care in the ICU was supportive. There were no deaths or serious complications in the study cohort. ❖

■ COMMENTARY BY DAVID J. PIERSON, MD

Angioedema produces asymmetrical, non-pitting edema of the skin and mucus membranes. Typically presenting with facial swelling, AE may progress to involve the tongue and supraglottic soft tissues, and fatal upper airway obstruction can ensue. Although it often is associated with allergy to foods (e.g., nuts or shellfish), stings by insects of the genus Hymenoptera, or administration of other medications (e.g., antibiotics), AE is increasingly encountered in patients taking ACEIs. It has been estimated that 0.1% to 0.2% of patients receiving ACEIs develop AE. With more than 40 million people

taking these agents in the United States today, this seemingly low incidence of a potentially life-threatening complication is important to clinicians working in EDs and ICUs.

The present study has a number of limitations. As with all retrospective studies, the true incidence of the entity being investigated cannot be determined. Cases of AE may have been miscoded or misdiagnosed. The association of AE with ACEI use may have been missed, and patients taking ACEIs may have had AE caused by something else. Clinicians should be cautious in generalizing these findings to other practice settings. For example, the finding that 91% of the patients in this series were African-American has uncertain implications; African-Americans compose 64% of inpatient admissions to the study hospital.

These cautions notwithstanding, this study is important in reminding us of a potentially life-threatening, but entirely treatable, complication whose frequency is substantial: At the study's 600-bed hospital, an average of nine patients with ACEI-induced AE were admitted each year during the study period.

Special Feature

Heparin Use in Acute Coronary Syndromes

By William J. Brady, MD

HEPARIN HAS BEEN SHOWN TO HAVE A PROFOUND synergistic effect with aspirin in preventing death, acute myocardial infarction (AMI), and refractory unstable angina. Heparin, either unfractionated or as the low-molecular-weight version, should be administered early in patients with AMI or the more severe forms of non-AMI acute coronary syndrome (ACS); patients with lower risk ACS presentations may not benefit from such therapy. The term heparin refers to a heterogenous collection of mucopolysaccharide chains of varying lengths and sizes. Unfractionated heparin (UFH) is composed of a mixture of polysaccharide chains with varying molecular weights. Low-molecular-weight heparin (LMWH) is approximately 30% of the molecular weight of UFH; the mucopolysaccharide chains of LMWH are more homogeneous in size. UFH binds to antithrombin III, forming a complex, which then inactivates both thrombin and activated factor X. The conversion to fibrin is prevented, ultimately inhibiting coagulation. UFH also potentiates the inactivation of factors XIa and IXa via antithrombin. Additional anticoagulant effects are mediated via platelet interaction. LMWH inhibits the coagulation cas-

cade in a similar fashion to UFH with action at varying levels of the cascade, including antithrombin III, thrombin, and factor Xa. It is theorized that LMWH exerts anticoagulant effects somewhat earlier in the coagulation cascade. All LMWHs are not necessarily equivalent in efficacy. The different forms of LMWH reportedly demonstrate variable antithrombotic effects attributed to differing ratios of anti-factors Xa to IIa.

Evidence-Based Support for Heparins in ACS

The FRISC trial¹ demonstrated that the combination of aspirin and LMWH significantly decreased the rate of non-fatal AMI or death at 1 week; this early beneficial effect was of a lesser magnitude later in the treatment course (40 to 150 days) coupled with an increased in minor bleeding. The FRIC study² reported the use of LMWH or UFH in patients with unstable angina and non-STEMI. The rates of recurrent ACS, urgent revascularization, and death were similar at 1 week and in 45 days.

In the ESSENCE study,³ a benefit was found with LMWH at 30 days coupled with an increased risk of minor bleeding. The ASSENT-3 study⁴ compared various combinations of LMWH (enoxaparin), UFH, and abciximab with tenecteplase in patients with AMI. Early in the course of therapy (48 hours), the LMWH group demonstrated a significantly lower rate of occurrence of death and recurrent ACS when compared with UFH infusion (6.1% vs 8.8%, respectively). At 30 days, the enoxaparin group continued to outperform UFH with a lower rate of occurrence of the primary endpoint when compared with UFH (15.4% vs 11.4%). The investigators concluded that LMWH was superior, particularly early in the treatment course, when compared with UFH therapy; efficacy was similar, however, at later stages of management. The investigators stated that "...in view of the present data and the ease of administration, enoxaparin might be considered an attractive alternative anticoagulant treatment when given in combination with tenecteplase."⁴

The TIMI-11B study⁵ compared LMWH (enoxaparin) with UFH in patients with unstable angina or non-STEMI. The patients managed with LMWH fared significantly better with fewer instances of recurrent AMI, need for urgent revascularization, or death. The initial benefit observed with LMWH was sustained through day 43; beyond this time, however, no further relative decrease in events was observed. In fact, there was an increase in the rate of major hemorrhage (spontaneous and instrumented) with chronic LMWH treatment.⁵

Comparing UFH and LMWH

In general, it appears that the LMWHs, particularly enoxaparin with its favorable anti-factor Xa/antifactor

IIa ratio, offers a short-term benefit in the unstable coronary patient, which lessens significantly beyond the first week of treatment. The advantages of LMWH over UFH are numerous. LMWH inactivates factor Xa, which is resistant to inactivation by UFH. LMWH has fewer binding characteristics to coagulation factors such as platelet factor 4, other plasma proteins, and endothelial cells, resulting in higher bioavailability. LMWH has a longer half-life and less individual variability of the anticoagulant response when compared with UFH. LMWH has a lower affinity for von Willebrand factor, increased vascular permeability, and a weak effect on platelet function. These differences could explain why LMWH produces less bleeding than UFH with equivalent or higher antithrombotic effects. The long half-life of LMWH and its predictable anticoagulant response to weight-adjusted doses allows once- or twice- daily subcutaneous administration without laboratory monitoring. LMWHs are the agents of choice when heparin is contraindicated by no availability of activated partial thromboplastin time (aPTT) measurements, but the disadvantage is expense.

Certainly, the majority of patients with AMI will require therapy with heparin, whether it be fractionated or unfractionated. Unstable angina, however, is an entirely different issue. Obviously, not all unstable angina cases are the same. For example, the stable patient with a classic description of new-onset angina who is sensation-free with a negative serum marker and a normal electrocardiogram (ECG) still is diagnosed correctly with unstable angina. Contrast this patient with the individual who presents with ongoing pain—either intermittent or constant—and a dynamic ECG; such a case clearly describes an active, unstable coronary event. The latter patient is in a higher-risk situation, and clearly will benefit from heparin therapy—more so than the patient in the former case. Heparin therapy is not without risk; it has been reported to be a major contributor to morbidity and mortality among hospitalized patients; it has been estimated that major bleeding will develop in one of every 90 patients treated and that one in 34 will develop heparin-induced thrombocytopenia. LMWH has been found to be as effective as UFH in patients with ACS and does not increase the bleeding risk while decreasing the risk of heparin-induced thrombocytopenia. Heparin should not be given to low-risk patients with unstable angina.⁶

Choosing the Dose of UFH and LMWH

In patients undergoing percutaneous coronary intervention, bleeding and mortality were higher in the TIMI 14 study⁷ in patients receiving 80 U/kg bolus and 18 U/kg/hr infusion compared with the 60 U/kg bolus and 16 U/kg/hr infusion. Therefore, the initial recommended

dose is 60 U/kg by intravenous bolus, followed by a maintenance infusion of 16 U/kg/hr. Heparin should be titrated so that the aPTT is 1.5 to 2.5 times control while using the maintenance infusion. UFH has some practical advantages over LMWH in the patient with AMI who is scheduled to undergo percutaneous coronary intervention in the immediate future; heparin infusions may be shut off should excessive bleeding complicate the situation, which is more likely as these patients receive glycoprotein IIb/IIIa inhibitors also. LMWH should be administered subcutaneously in a twice-daily regimen at a dose of 1 mg/kg; doses should be reduced for patients with pronounced renal insufficiency. Absolute contraindications to heparin therapy include known allergy, history of heparin-induced thrombocytopenia (only for UFH), active ongoing life-threatening hemorrhage, or predisposition to such hemorrhage. ❖

References

1. Wallentin L, et al for the FRISC group. Low molecular weight heparin during instability in coronary artery disease. Fragmin during instability in coronary artery disease (FRISC) study group. *Lancet* 1996;347:561-568.
2. Klein W, et al. Comparison of low-molecular-weight heparin with unfractionated heparin acutely and with placebo for 6 weeks in the management of unstable coronary artery disease. Fragmin in unstable coronary artery disease study (FRIC). *Circulation* 1997;96:61-68.
3. Cohen M, et al. A comparison of low-molecular-weight heparin with unfractionated heparin for unstable coronary artery disease. Efficacy and Safety of Subcutaneous Enoxaparin in Non-Q-Wave Coronary Events Study Group. *N Engl J Med* 1997;337:447-452.
4. Assessment of the Safety and Efficacy of a New Thrombolytic Regimen (ASSENT)-3 Investigators. Efficacy and safety of tenecteplase in combination with enoxaparin, abciximab, or unfractionated heparin. The ASSENT-3 randomised trial in acute myocardial infarction. *Lancet* 2001;358:605-613.
5. Antman EM, et al. Enoxaparin prevents death and cardiac ischemic events in unstable angina/non-Q-wave myocardial infarction. Results of the thromboly-

sis in myocardial infarction (TIMI) 11B trial. *Circulation* 1999;100:1593.

6. Brewster GS, et al. Heparin should be administered to every patient admitted to the hospital with possible unstable angina. *West J Med* 2000;173:138-140.
7. Schweiger MJ, et al. Early coronary intervention following pharmacologic therapy for acute myocardial infarction (the combined TIMI 10B-TIMI 14 experience). *Am J Cardiol*. 2001;88:831.

Physician CME Questions

1. **One advantage of low-molecular-weight heparin over unfractionated heparin is:**
 - a. there is no need to monitor the activated partial thromboplastin time.
 - b. it can be shut off if excessive bleeding occurs.
 - c. its dosing is not affected by renal insufficiency.
 - d. its dosing is not weight-based.
2. **The highest rates of survival after ED thoracotomy are reported among patients with:**
 - a. stab wounds to the neck.
 - b. gunshot wounds to the chest.
 - c. stab wounds to the chest.
3. **Compared with regular insulin IV therapy, treatment of diabetic ketoacidosis (DKA) with insulin lispro by SQ injection:**
 - a. results in comparable rates of hypoglycemic events.
 - b. requires less frequent monitoring of blood glucose.
 - c. reduces length of hospital stay.
 - d. results in slower resolution of ketoacidosis.
4. **In the report by Hsieh et al, ultrasound confirmation of endotracheal tube placement relied on visualizing:**
 - a. the upper airway and endotracheal tube location.
 - b. appropriate diaphragm motion with ventilation.
 - c. appropriate hepatic and gastric movement with ventilation.
 - d. the esophagus without the endotracheal tube.

Answers: 1.a; 2.c; 3.a; 4.b.

CME Objectives

To help physicians:

- Summarize the most recent significant emergency medicine-related studies;
- Discuss up-to-date information on all aspects of emergency medicine, including new drugs, techniques, equipment, trials, studies, books, teaching aids, and other information pertinent to emergency department care; and
- Evaluate the credibility of published data and recommendations.

CME Instructions

Physicians participate in this continuing medical education program by reading the article, using the provided references for further research, and studying the questions at the end of the article. Participants should select what they believe to be the correct answers, then refer to the list of correct answers to test their knowledge.

To clarify confusion surrounding any questions answered incorrectly, please consult the source material. After completing this activity, you must complete the evaluation form that will be provided at the end of the semester and return it in the reply envelope provided to receive a certificate of completion. When your evaluation is received, a certificate will be mailed to you.

ECG Review

A Three-Beat Run in Lead V₁

By Ken Grauer, MD




Figure. 12-lead ECG obtained from a 57-year-old woman with palpitations. Note the three-beat run in lead V₁.

Clinical Scenario: The electrocardiogram (ECG) in the Figure was obtained from a 57-year-old woman with palpitations. Is there a short run of ventricular tachycardia (VT) in lead V₁? What else may be wrong with the tracing?

Interpretation: The rhythm is rapid and irregularly irregular. The absence of P waves defines this as atrial fibrillation with a rapid ventricular response. Three abnormal looking beats are seen consecutively in lead V₁ (beginning with beat Y). Although at first glance one might be tempted to interpret this three-beat run as a salvo of VT, it is much more likely that these three beats are supraventricular in etiology and conducted with aberration. Each of these three beats manifests an rsR' morphology consistent with a right bundle-branch block pattern (RBBB). Note that the initial direction and magnitude of the small r wave is identical for the three anomalous and the three narrow beats that occur in this lead. This is consistent with the conduction defect that occurs with RBBB, in which the initial vector of left-to-right septal depolarization is unaffected by the conduction disturbance. In further support that these beats are aberrantly conducted is the finding of a reason for aberrant conduction: the *coupling* interval of the first anomalous complex (the distance between beat X and beat Y) is short (increased likelihood that there will be relative refractori-

ness of the ventricular conduction system). In addition, the preceding R-R interval (i.e., the R-R interval before beat X) is relatively long. Because the length of the refractory period is determined by the duration of the *preceding* R-R interval, the opportunity for aberrant conduction to occur is enhanced when early occurring beats (such as Y) are preceded by a relatively longer R-R interval. Finally, the complex labeled Z (in lead V₆) manifests a QRS consistent with incomplete RBBB conduction (it is the only complex in lead V₆ with an S wave), thereby providing additional evidence of RBBB-type conduction delay for selected beats on this tracing. Although not infallible, the combination of morphologic features described above (that are consistent with a bundle-branch block form of conduction delay) occurring in a setting that predisposes to aberrant conduction (i.e., rapid atrial fibrillation) strongly suggest that the three-beat run seen in lead V₁ is not VT. Rate control of this patient's rapid atrial fibrillation probably will result in resolution of anomalous complexes. Otherwise, there is low-voltage, non-specific ST-T wave abnormalities and a highly unusual negative complex in lead I. In view of the upright QRS complex in lead aVR, limb lead reversal would not negate our interpretation of the rhythm, nor of the high likelihood for aberrant conduction of the three-beat run that is seen in the precordial leads. ♦

In Future Issues:

Clinical Characteristics of Adult Meningitis

Trauma Reports

Vol. 5, No. 6

Supplement to *Emergency Medicine Reports, Pediatric Emergency Medicine Reports, ED Management, and Emergency Medicine Alert*

Nov./Dec. 2004

As an isolated injury or a frequent component in a multiple trauma patient, the knee is one of the most commonly injured joints. Although not usually a life-threatening injury, correct management may significantly affect the quality of life and mobility for a patient following a traumatic event. In addition, failure to make a timely and accurate diagnosis of a vascular injury may have significant detrimental effects for the trauma patient and result in permanent disability.

It is critical that emergency department (ED) physicians use the history of the injury to direct the physical examination and diagnostic testing, as well as provide prompt referral to an orthopedic specialist if the condition requires surgical intervention. The ED physician also must understand the limitations faced in making a definitive diagnosis in the first 24 hours following an acute injury, a critical time period.

The authors present a thorough review of the anatomy, critical features of the physical examination, and indications for further diagnostic testing in a patient who has sustained a traumatic knee injury.

— The Editor

Introduction

Traumatic knee injuries are a common reason for presentation to EDs following injury; approximately 1 million knee injuries are treated in EDs in the United States each year.^{1,2} In addition to isolated knee injuries, the knee joint is frequently injured in association with other injuries in multiple trauma victims.

Because the knee is a complex joint, it is critical to understand the different patterns of injury that can occur: injuries to bone, ligaments, cartilage, and surrounding soft tissue.

The emergency physician (EP) must develop a rational imaging strategy to identify potential injuries and have a good understanding of the types of injuries to properly care for the patient with knee trauma.

Many traumatic knee injuries require orthopedic consultation for operative management, and those injuries must be identified quickly and efficiently to expedite care of the patient.

Finally, there are many pitfalls in the evaluation of the knee, especially the failure to diagnose arterial injury, which makes a thorough understanding of the knee and its surrounding structures essential to the active clinician.

Knee Trauma: Assessment, Diagnostic Evaluation, and Management

Authors: **Howard A. Werman, MD**, Professor of Clinical Emergency Medicine, The Ohio State University College of Medicine and Public Health; Medical Director, MedFlight, Columbus, OH; **Robert E. Paasche, MD**, Emergency Medicine Physician, Emergency Physicians, PC, Benefis Hospital, Great Falls, MT; **M. Jake Ott, III, MD**, Attending Physician, St. Ann's Medical Center, Columbus, OH.

Peer Reviewer: **George L. Foltin, MD, FAAP, FACEP**, Director, Pediatric Emergency Medicine, Bellevue Hospital Center/New York University Medical Center, New York.

Now available online at www.ahcpub.com/online.html or call (800) 688-2421 for more information.

EDITOR IN CHIEF

Ann Dietrich, MD, FAAP, FACEP
Associate Clinical Professor
Ohio State University
Attending Physician
Columbus Children's Hospital
Associate Pediatric Medical Director
MedFlight
Columbus, Ohio

EDITORIAL BOARD

Mary Jo Bowman, MD
Associate Professor of Clinical Pediatrics
Ohio State University College of Medicine
Attending Physician, Children's Hospital of Columbus
Columbus, Ohio

Larry N. Diebel, MD
Associate Professor of Surgery
Detroit Medical Center
Wayne State University
Detroit, Michigan

Robert Falcone, MD
Senior Operations Officer
Grant Medical Center
Columbus, Ohio

Theresa Rodier Finerty, RN, MS
Director, Emergency and Trauma Services,
OSF Saint Francis Medical Center
Peoria, IL

Dennis Hanlon, MD
Director
Emergency Medicine Residency Program
Assistant Professor of Emergency Medicine
Allegheny General Hospital
Pittsburgh, Pennsylvania

Robert Jones, DO, FACEP
Emergency Ultrasound Coordinator
OUCOM/Doctor's Hospital Emergency Medicine
Residency Program
Columbus, Ohio
Attending Physician, MetroHealth Medical Center
Cleveland, Ohio

S.V. Mahadevan, MD, FACEP
Assistant Professor of Surgery
Associate Chief, Division of Emergency Medicine
Stanford University School of Medicine
Stanford, California

Ronald M. Perkin, MD, MA, FAAP, FCCM
Professor and Chairman
Department of Pediatrics
Brody School of Medicine at East Carolina University
Medical Director, Children's Hospital University
Health Systems of Eastern Carolina
Greenville, North Carolina

Steven A. Santanello, DO
Medical Director, Trauma Services
Grant Medical Center
Columbus, Ohio

Eric Savitsky, MD
Assistant Professor of Medicine
Emergency Medicine/Pediatric Emergency Medicine
UCLA Emergency Medicine Residency Program
Los Angeles, California

Perry W. Stafford, MD, FACS, FAAP, FCCM
Head, Pediatric Surgery
Jersey City Medical Center
Jersey City, New Jersey

© 2004 Thomson American Health Consultants
All rights reserved

Anatomy of the Knee

An understanding of the anatomy of the knee is imperative to assessing injuries involving this complex joint. The knee is the largest articular joint in the body and consists of a hinge joint formed by the articulation of the femur and the tibia. The femur broadens out at the knee joint into two rounded articular prominences: the medial and lateral condyles. Those prominences articulate with the tibial plateaus on the proximal tibia. Small eminences from the condyles form the medial and lateral epicondyles, which serve as attachment points for the collateral ligaments. The anterior portion between both condyles contains a shallow depression that accommodates the patella.

The patello-femoral articulation is not involved in weight bearing.² The proximal tibia also contains medial and lateral condyles, which are flattened surfaces that support weight bearing. These surfaces are referred to as the medial and lateral tibial plateaus. Each plateau has approximately 10° of downward slope away from the center of the tibia. Centrally, the tibial plateau

Trauma Reports™ (ISSN 1531-1082) is published bimonthly by Thomson American Health Consultants, 3525 Piedmont Road, N.E., Six Piedmont Center, Suite 400, Atlanta, GA 30305. Telephone: (800) 688-2421 or (404) 262-7436.

Vice President/Group Publisher: Brenda Mooney

Editorial Group Head: Valerie Loner

Managing Editor: Martha Jo Dendinger

Marketing Manager: Schandale Kornegay

POSTMASTER: Send address changes to **Trauma Reports**, P.O. Box 740059, Atlanta, GA 30374. Copyright © 2004 by Thomson American Health Consultants, Atlanta, GA. All rights reserved. Reproduction, distribution, or translation without express written permission is strictly prohibited.

Accreditation

Trauma Reports™ continuing education materials are sponsored and supervised by Thomson American Health Consultants. Thomson American Health Consultants designates this continuing education activity for up to 2.5 hours in Category 1 credit toward the AMA Physician's Recognition Award. Each physician should claim only those hours of credit that he/she actually spent in the educational activity. This CME activity was planned and produced in accordance with the ACCME Essentials. Approved by the American College of Emergency Physicians for 2.5 hours of CEP Category 1 credit.

Thomson American Health Consultants (AHC) is accredited by the Accreditation Council for Continuing Medical Education (ACCME) to provide continuing medical education for physicians.

Trauma Reports® is approved for approximately 2.5 nursing contact hours. This offering is sponsored by Thomson American Health Consultants, which is accredited as a provider of continuing education in nursing by the American Nurses'

THOMSON
★
**AMERICAN HEALTH
CONSULTANTS**

Conflict of Interest Disclosure

In order to reveal any potential bias in this publication, and in accordance with Accreditation Council for Continuing Medical Education guidelines, Dr. Dietrich (editor in chief), and editorial board members Bowman, Diebel, Falcone, Finerty Hanlon, Jones, Mahadevan, Perkin, Santanello, Savitsky, and Stafford report no relationships with companies related to the field of study covered by this CME program. Drs. Werman, Paasche, and Ott (authors) and Foltin (peer reviewer) report no relationships with companies related to the field of study covered by this CME program.

Subscriber Information

Customer Service: 1-800-688-2421

Customer Service E-Mail: customerservice@ahcpub.com

Editorial E-Mail: martha.dendinger@thomson.com

World Wide Web page: <http://www.ahcpub.com>

FREE to subscribers of *Emergency Medicine Reports*, *Pediatric Emergency Medicine Reports*, *Emergency Medicine Alert*, and *ED Management*.

Credentialing Center's Commission on Accreditation. Provider approved by the California Board of Registered Nursing, Provider Number CEP 10864, for approximately 2.5 contact hours. This program (#0105-1) has been approved by an AACN Certification Corp.-approved provider (#10852) under established AACN Certification Corp. guidelines for 2.5 contact hours, CERP Category A.

This is an educational publication designed to present scientific information and opinion to health professionals, to stimulate thought, and further investigation. It does not provide advice regarding medical diagnosis or treatment for any individual case. It is not intended for use by the layman. Opinions expressed are not necessarily those of this publication. Mention of products or services does not constitute endorsement. Clinical, legal, tax, and other comments are offered for general guidance only; professional counsel should be sought for specific situations.

This CME/CE activity is intended for emergency, family, osteopathic, trauma, surgical, and general practice physicians and nurses who have contact with trauma patients.

It is in effect for 36 months from the date of publication.

For Customer Service,

Please call our customer service department at (800) 688-2421. For editorial questions or comments, please contact **Martha Jo Dendinger**, Managing Editor, at martha.dendinger@thomson.com.

contains two superior projections: the anterior and posterior tibial spine. These projections serve as attachment points for the cruciate ligaments.

Also of interest is the tibial tuberosity, which is located on the anterior surface of the tibia below the knee. This bony prominence serves as the attachment of the quadriceps tendon. The proximal fibula, although not technically part of the knee, is included in this discussion. These structures combine to form the major bony articulations of the knee. (See *Figure 1*.)


The popliteal fossa is a depression found behind the knee joint. Several important structures are located in this recess, including the popliteal artery and vein, as well as the peroneal and tibial nerves. The popliteal artery represents the continuation of the femoral artery into the knee joint. This artery supplies a rich anastomosis of blood vessels supplying the knee joint and then divides into the anterior and posterior tibial arteries. The artery is relatively fixed within the popliteal fossa, thus, making it prone to injury with any stress on the knee joint.^{3,4}

The common peroneal nerve wraps around the proximal fibula. Injury to this nerve may result in a foot drop.⁵ Intact dorsiflexion at the ankle must be documented with any significant trauma to the proximal fibula.

The static stability of the knee joint depends on its surrounding ligaments (See *Figure 1*). The anterior cruciate ligament (ACL) prevents anterior displacement of the tibia in relation to the femur. The posterior cruciate ligament (PCL) prevents posterior and rotational displacement. Both the ACL and PCL also contribute to proprioception. The medial collateral ligament (MCL) resists valgus stress, while the lateral collateral ligament (LCL) resists varus stress.


With the knee in full extension, all of the ligaments are taut, ensuring complete stability of the knee joint. The knee must be flexed to 20-40° before the joint is relaxed, and some rotational movement of the joint is possible. The medial and lateral menisci are cartilaginous structures that support the femorotibial articulation and also aid in weight bearing. (See *Figure 2*.) The menisci contribute to joint stability and provide both lubrication and cushioning to the knee. The medial meniscus is attached firmly to the MCL and joint capsule making it more susceptible to injury. Because of the absence of pain fibers, menisci injuries may have a delayed detection or go undetected.

The muscular structures of the knee provide dynamic stability to the joint. The knee extensors (quadriceps group) include the vastus medialis, vastus intermedius, vastus lateralis, and rectus femoris. They combine to form the quadriceps tendon, which inserts into and encompasses the patella. The quadriceps tendon then becomes the patellar tendon and inserts into the tibial tuberosity. The vastus medialis exerts a medial force on the patella, thus, preventing lateral subluxation. The knee flexors (hamstrings group) include the semimembranosus, semitendinosus, and biceps femoris. The knee adductors include the gracilis, sartorius, and a portion of the semitendinosus, combining to form the pes anserinus medially. The iliotibial tract, popliteus muscle, and a portion of the biceps femoris attach laterally.

Figure 1. Knee Bony Anatomy

The bony anatomy of the knee consists of the articulation of the femur and the tibia, which meet to form a hinge joint. The joint is protected in front by the patella. The joint is cushioned by articular cartilage that is located at the ends of the tibia, femur, and under the patella. The lateral and medial meniscus are cartilaginous and further cushion the joint.

Reprinted with permission from: The Nicholas Institute of Sports Medicine and Athletic Trauma. Available at: www.nis-mat.org.

Figure 2. Knee Ligamentous Anatomy

The ligaments of the knee are designed to stabilize the knee. The collateral ligaments are located along the sides of the knee and limit lateral motion. The ACL is located at the center of the knee and connects the tibia to the femur. The ACL functions to limit rotation and forward motion of the tibia. The PCL limits backward motion of the tibia. The MCL restricts valgus stress, and the LCL resists varus stress.

Reprinted with permission from: The Nicholas Institute of Sports Medicine and Athletic Trauma. Available at www.nis-mat.org.

History and Physical Examination

A thorough history regarding the mechanism of injury is crucial to identification of the particular pattern of knee injury. For example, an anterior force applied to the tibia with the foot planted typically results in an ACL injury; the PCL frequently is damaged by a posterior force under the same conditions.^{6,7}

It should be noted, however, that an accurate history of the injury may be absent in patients with multiple trauma; therefore, a careful examination of the knee is imperative and correlation with historical information appropriate in all patients. Information obtained should include the direction of the force that caused the injury and the weight-bearing status of the patient at the time of impact. The knee is commonly subjected to compression, distraction, flexion, extension, rotation, valgus or varus stress, and sliding forces.

Additional information, including the position of the patient in an automobile (e.g., driver, front-seat or rear-seat passenger), anticipated impact, and damage to the dashboard, may be particularly important in association with motor vehicle trauma. The location of pain and the presence of a pop or snap (noted in up to 90% of cases associated with ACL rupture at the time of injury) are also very important.⁸⁻¹⁰

Patients with ACL or PCL injuries also may complain of the knee “giving out,” especially with pivoting. Complaints of knee

locking may indicate a torn meniscus, a loose body, rupture of the ACL or PCL, or an osteochondral fracture.¹¹⁻¹²

An osteochondral fracture occurs when a portion of the articular surface, including both lining cartilage and underlying bone, becomes separated from either the femur or tibia. The time of injury and the amount of swelling or effusion that has developed can be helpful in distinguishing the injured structure. The cruciate ligaments are well vascularized, and a tear can produce a large effusion within a couple of hours.^{8,9,13,14} Acute hemarthroses is a sign of intra-articular injury, with a high percentage of these injuries involving ACL tears. Osteochondral fractures also can produce large effusions quickly and account for as many as 67% of traumatic hemarthroses in children without apparent fracture on x-ray.¹⁵ A smaller effusion that collects during a longer period of time is more indicative of a partial ligament tear or meniscal injury.

A detailed examination of the knee in flexion and extension can provide very specific information about the type of injury involved. The initial evaluation should include assessment of deformity, swelling, effusion, tenderness, and open wounds. It is also helpful to compare any findings to the uninjured knee. Location of tenderness on examination corresponds with the site of injury identified surgically in up to 76% of cases.¹⁰ As noted above, the presence of a large effusion that is noted within 1-2

Figure. 3 The Lachman Test



The Lachman's test is more sensitive than the anterior drawer test for ACL rupture. As illustrated above, the knee should be flexed 20-30°. One hand should hold the distal femur. With the other hand, apply anterior force to the proximal tibia. If an end-point is not appreciated or significant movement of the anterior tibia occurs, the test is considered positive.

Reprinted with permission from The Nicholas Institute of Sports Medicine and Athletic Trauma. Available at www.nismat.org.

hours of injury is highly suggestive of cruciate ligament rupture or an osteochondral fracture. A full popliteal fossa may be seen with popliteal artery disruption, suggesting that the knee may have been dislocated, that a bicruciate ligament rupture may be present, or that both injuries have occurred simultaneously.

It is extremely important to remember that a reliable examination may not be possible within the first few hours of injury, and the patient may need to have a repeat exam in 24-48 hours when the pain and swelling have subsided. In the multiply injured patient, other priorities may limit the initial examination of the knee and associated structures. Following stabilization, a complete and directed physical examination will reveal likely injury patterns.¹⁶

Valgus and varus stress should be applied to the knee in 20-30° of flexion and in full extension. This action will help to identify MCL and LCL injuries. Examination of the knee in slight flexion allows for relaxation of the major muscle groups that provide support to the knee and isolates the supporting ligamentous structures of the knee joint.

Anterior and posterior drawer tests should be performed with the hip flexed at 45°, the knee flexed at 90°, and the foot secured in a neutral position on the exam table. Anterior force should be applied to the proximal tibia to assess for ACL integrity and posterior force applied to assess for PCL integrity. The same examinations also should be repeated with the foot secured in internal and external rotation.² The tibia should be limited by a discrete end-point in the anterior and posterior directions. Any sliding

Figure 4. The McMurray Test



The examiner flexes the knee beyond 90° with the foot externally rotated in the examiner's hand. The other hand palpates the medial joint line as the knee is extended gradually beyond 90°. A medial meniscus injury is indicated by pain, a popping sensation, or crepitus.

Reprinted with permission from The Nicholas Institute of Sports Medicine and Athletic Trauma. Available at www.nismat.org.

forward or backward suggests an ACL or PCL injury.

Lachman's test is more sensitive for ACL rupture (up to 99% sensitivity vs 77% for the anterior drawer test).¹⁷ Hold the distal femur with one hand, and apply anterior force to the proximal tibia with the other hand while the knee is flexed 20-30° (*See Figure 3.*). Here again, a discrete end-point should be appreciated. A lack of a discrete end-point or significant anterior movement of the tibia is indicative of a positive test.

The Lateral Pivot Shift test is performed with the knee in full extension. The leg is lifted by the distal tibia, which will produce subluxation of the proximal tibia with ACL disruption. The knee is flexed carefully to 20-40°, while a mild valgus force is applied to the knee, and the foot is moved laterally. A thud can be appreciated as the tibia subluxes against the femur with this motion.

Meniscal injuries are detected using the McMurray's test, Apley compression test, or medial-lateral grind test.¹⁸ The McMurray test involves flexion of the knee beyond 90° with the foot externally rotated in the examiner's hand. (*See Figure 4.*) The other hand palpates the medial joint line as the knee is extended gradually beyond 90°. Pain, a popping sensation, or crepitus indicates a medial meniscus injury. Repeat the test with the foot internally rotated while the lateral joint line is palpated. Pain and crepitus with flexion indicate lateral meniscus injury. The McMurray's test alone has been shown to have a low sensitivity and specificity for detecting medial meniscus injuries.^{19,20}

Apley's compression test is performed with the patient in a prone position, and the knee flexed to 90°. Pain with rotation of

Figure 5. Intercondylar Femoral Fracture



This patient sustained an intercondylar femur fracture. Because a large amount of force is necessary to produce this injury, the patient should be assessed carefully for other injuries. In addition, the close proximity of the popliteal artery and vein to the distal femur should alert the clinician to the possibility of vascular injury.

Image courtesy of Howard Werman, MD.

the foot while upward traction is applied suggests damage to the capsule and/or ligaments. Pain with rotation of the foot while downward traction is applied indicates meniscal injury.

The medial-lateral grind test is performed in the supine patient by cradling the calf in one hand and palpating the medial and lateral joint lines with the other. Valgus and varus stress is applied to the joint. Any crepitus felt along the joint line indicates possible meniscal injury.

Bragard's sign is tenderness along the anterior medial joint line upon internal rotation and extension of the tibia and indicates a likely medial meniscus injury.

Payr's sign is pain with pressure applied to the medial aspect of the knee with the patient sitting cross-legged. The knees should be flexed at 90°. This finding is also consistent with medial meniscus injury.

First Steinmann's sign is either pain in the anterolateral joint space with internal rotation and flexion of the knee, which is consistent with a lateral meniscus tear, or pain in the anteromedial joint space with external rotation and flexion of the knee, which suggests a medial meniscus tear.

Secondary Steinmann's sign helps to distinguish a meniscal injury from an LCL or MCL disruption. It involves a shifting

Figure 6. Tibial Plateau Fracture



This patient sustained a tibial plateau fracture. Meniscal and ligamentous injuries commonly accompany these fractures.

Image courtesy of Howard Werman, MD.

location of tenderness from the region of the collateral ligament when the knee is flexed to the anterior aspect of the knee when fully extended.

In addition, a thorough evaluation of the neurovascular status of the knee should be done, focusing on the presence or absence of pulses in the posterior tibial, dorsalis pedis, and popliteal arteries, as well as the sensory function of the various nerves supplying the area below the knee.

Injury to the Bony Structures of the Knee

Bony injuries to the knee can involve the distal femur, proximal tibia and fibula, patella, or dislocation of the knee. The management of these various injuries differs based on the mechanism and location of injury.

Femoral Fractures. Fractures of the distal femur are rare injuries^{1,21,22} and are grouped into four different categories: Class A, supracondylar fractures; Class B, intercondylar fractures; Class C, condylar fractures; and Class D, epiphyseal fractures (See Figure 5.)

All femoral fractures present with knee pain or hip pain, inability to bear weight, and tenderness over the distal femur. These fractures usually occur as the result of a direct blow to the distal femur, such as from a dashboard or steering wheel. A large amount of force is required to produce a femoral fracture and

consequently, these fractures often are associated with other injuries. Additionally, because of the muscular attachments of the quadriceps group, the hamstring muscles and both the gastrocnemius and soleus muscles, displacement of the fracture fragments is common. Remember that the popliteal artery and vein are in close proximity to the distal femur and are prone to injury with such fractures.

Supracondylar Fractures. By definition, these fractures occur above the femoral condyles and do not involve the knee joint. As a result, effusions are not associated with this type of injury. Five percent to 10% of these fractures are open fractures.²³ All of the other fractures mentioned above have an intra-articular component. Epiphyseal fractures usually occur in children older than 10 years. Because bone growth around the knee joint accounts for almost two-thirds of adult leg length, these injuries commonly are associated with leg shortening, and all Salter-Harris fractures of the knee joint should have urgent orthopedic evaluation.^{10,21,22}

Distal femur fractures all require urgent orthopedic consultation and can be treated with immobilization, traction, or operative fixation at the discretion of the consulting orthopedic specialist. Skeletal traction often is the preferred method by orthopedic surgeons once the patient is admitted; however, a hare traction splint can be used to immobilize the fracture in the ED until the skeletal pins can be placed. Significant pain control often is required; traction on the femur is extremely painful.² Commonly associated injuries include ipsilateral hip fractures, popliteal vascular injuries, peroneal nerve injury, and damage to the quadriceps or hamstring groups.

Tibial Fractures. Tibial fractures are divided into four classes: Class A, condylar fractures; Class B, spine fractures; Class C, tuberosity fractures; and Class D, subcondylar fractures.

Tibial fractures typically occur in one of two ways: 1) a direct blow to the tibia (bumper injury); or 2) axial compression of the tibia, which occurs when the knee is forcibly extended in anticipation of a frontal impact.

Class A injuries (condylar and plateau fractures) usually are produced by axial loading (e.g., a fall from a height). (See Figure 6.) The rotational stress on the knee will determine whether the medial or lateral tibial plateau is affected.²⁴ The lateral plateau is the site of fracture in the majority of cases.¹ Meniscal and ligamentous injuries commonly accompany these fractures; the menisci and medial and lateral collateral ligaments are related closely to and attached to the tibial plateaus.²⁵ A plateau fracture without depression of the articular surface can be treated by a long leg cast. Any depression or irregularity of the articular surfaces requires immediate orthopedic consultation and surgical repair. ACL and MCL tears are associated with lateral plateau fractures; PCL and LCL tears are associated with medial plateau fractures. Lateral plateau injuries are more common and usually are due to involvement of the tibial spine.²⁵

Class B injuries (spine fractures) are rare but present similar to ACL and PCL tears, depending upon which spine is fractured. The mechanism of injury typically involves an anterior or posterior force to the proximal tibia with the knee in flexion. Anterior

Figure 7. Patellar Fracture



This radiograph demonstrates a patellar fracture. Patellar fractures typically occur secondary to direct knee trauma. Non-displaced patellar fractures may be managed with a knee immobilizer and pain control.

Image courtesy of Howard Werman, MD.

spine fractures are 10 times more common than posterior spine fractures and are associated with ACL injuries.²⁵ Complete spine avulsion requires urgent orthopedic consultation and surgical repair, while incomplete fractures can be treated by a knee immobilizer and orthopedic follow-up.

Class C injuries (tuberosity fractures) involve the attachment of the quadriceps tendon, and present with pain over the anterior tibia with signs similar to either partial or complete quadriceps tendon rupture (difficulty or inability to perform a straight leg raise). These fractures can occur as the result of a shearing force or more commonly, due to forced flexion of the knee, creating stress on the quadriceps tendon. Partial avulsions can be treated by immobilization and orthopedic follow-up; complete avulsions require operative fixation. Clinically, this type of injury is indistinguishable from quadriceps tendon rupture, but plain x-rays

Figure 8. Angiogram of Popliteal Artery Showing Intimal Tear



Injury to the popliteal artery is common in patients with knee dislocations. These injuries require early recognition and, optimally, a repair within 6-8 hours of the injury. Intact pulses in the foot do not exclude vascular injury.

Image courtesy of Howard Werman, MD.

will show the avulsed segment of the tibial spine attached to the end of the quadriceps tendon.

Class D injuries (subcondylar fractures) can be treated with immobilization and do not require urgent orthopedic consultation unless there is comminution of the fragments or associated injuries. Like supracondylar patellar fractures, these injuries do not involve the joint space.

Proximal Fibular Fractures. These fractures are typically noted as the result of a direct blow to the area or a valgus stress on the knee joint. The latter mechanism typically produces lateral collateral ligament injury and peroneal nerve injury.¹ Isolated proximal fibular fractures can be treated symptomatically with a

knee immobilizer and analgesics. A variant of the proximal fibular fracture, the Maisonneuve's fracture, involves a distal tibial fracture or ligamentous disruption, resulting in instability of the ankle. Thus, the ankle joint should be assessed carefully if a proximal fibular fracture is present, and x-rays should be obtained if the ankle is tender.

Patellar Fractures. Patellar fractures typically are caused by direct knee trauma (e.g., a knee striking a dashboard), and are divided into several classes: non-displaced, transverse, pole fractures, comminuted (non-displaced or displaced), vertical, or osteochondral. High impact injuries are associated with hip fractures and dislocations. A secondary mechanism may involve an avulsion caused by the pull of the quadriceps mechanism. Patellar fractures are more common in males.

A non-displaced fracture can be treated with a knee immobilizer and pain control. If the quadriceps mechanism is intact (See Figure 7), the patient can bear weight as well. Fractures associated with disruption of the extensor mechanism or displacement more than 3 mm requires operative repair in an urgent manner.

Remember that anatomical variants of the patella can include a bipartite and tripartite patella. With a bipartite patella, a secondary ossification center typically is noted in the upper outer portion of the patella, which may be confused with a fracture. Because these variants tend to be bilateral, a comparison view of the uninjured patella should be obtained.

Patellar Dislocation

Patellar dislocations typically occur when the patient twists on an extended knee. Rarely, dislocation can occur as the result of a lateral or medial blow to the patella. The condition is more common in women.²⁶ Many patients (10–50%) will have recurrent episodes during their lifetimes. The patella almost always dislocates laterally (80%).²⁶ Sometimes this injury is associated with medial joint capsule injury.

Patellar dislocation is very painful, and the patient often presents with a fixed, partially flexed knee with obvious displacement of the kneecap. The reduction can be performed in one of two manners.

Passive Hyperextension of the Knee. This is a painful procedure and often requires pain control and sedation. The lower leg is grasped in one hand, the upper leg in the other, and the knee is straightened passively. The patella then is moved back into place by hand, forcing the patella medially.

Active Hyperextension of the Knee. The patient can be asked to straighten the knee (e.g., to touch the bed with the back of his knee while standing or lying). A cooperative patient can perform this maneuver, and reduction causes immediate cessation of pain.

Whichever method is chosen, plain knee radiographs should be obtained following successful reduction to identify any associated patellar fractures or osteochondral injuries, which are seen in up to 5% of cases.^{1,26} Additionally, a thorough knee examination should be performed once the patella is in anatomic alignment; up to 12% of patients with a patellar dislocation have an associated ligamentous or meniscal injury.^{1,26} Knee immobiliza-

tion should be used to prevent recurrent dislocation, and follow-up within one to two weeks is indicated. If recurrent dislocation occurs, operative fixation may be needed.

Knee Dislocation


Knee dislocation is the result of a significant force delivered to the knee and occurs as the result of hyperextension, posterior forces to the tibia, or significant rotary forces.²⁷ One example is a bumper impact by a motor vehicle against a planted lower extremity. Any knee dislocation is associated with significant ligamentous disruption.⁵ There are three types of dislocation of the knee: anterior, posterior, and lateral. In anterior dislocations (the most common type), the tibia is displaced anteriorly to the femur and typically is the result of hyperextension of the knee joint. In posterior dislocations, the tibia is forced posteriorly by a direct force applied to the proximal tibia. Lateral dislocations can be medial or lateral, and occur with equal frequency. Lateral dislocations can be complicated further with rotary deformity of the knee as well. Any type of knee dislocation carries a 20% incidence of open joint injury⁵ and classically presents with a deformed knee and severe pain. Pulses in the foot must be assessed immediately and may be present or absent. Reduction of all three types of dislocations requires longitudinal traction. Sedation and pain relief commonly are required.^{1,5} Post-reduction radiographs should be obtained to assess for concomitant fractures, which can include tibial and patellar fractures, as well as fibular head fractures. A thorough neurovascular examination must be conducted following successful reduction to determine if there is injury to the peroneal and tibial nerves. One additional injury pattern is worthy of mention—proximal tibiofibular dislocations or subluxations—rare injuries that are seen after a significant fall. Peroneal nerve injuries commonly are associated with this type of dislocation.

Injury to the popliteal artery is common in knee dislocations (up to 40% of all dislocations).^{3,28} (See Figure 8.) It is critical that such injuries be identified early and repaired within 6-8 hours of injury.²⁸ Intact pulses in the foot do not exclude the possibility of arterial injury. An arteriogram should be performed on these patients according to most experts, although an alternative strategy of admission and repeated clinical examination after normal ankle brachial indices (ABI) in the ED has been suggested.⁵ An ABI is performed by obtaining a blood pressure in the arm, by standard technique, then taking a blood pressure at the ankle. Dividing the ankle pressure by the arm pressure determines the ABI. Ankle pressure should be greater than brachial pressure. Thus, a ratio of one or greater is considered normal. An ABI of 0.9 or less is concern for possible arterial injury. Patients who have peripheral arterial disease also will have an ABI of less than 0.9, possibly leading to a false positive result. Whichever strategy is used, knee immobilization and urgent orthopedic consult is required.

Treatment of Bony Knee Injuries

Many bony knee injuries can be treated by casting; immobilization of the joint above and below the injury is critical for opti-

Figure 9. Knee Ligamentous Injury



Sprains involve injury to the ligamentous structures of the knee. They vary in severity from a Grade 1 sprain (a small incomplete with no instability of the knee) to a Grade 3 sprain (a complete ligamentous disruption with complete joint instability).

Reprinted with permission from Hughston Sports Medicine Foundation.

healing. For tibial and fibular fractures, it can be accomplished either with a long leg cast or through the combined use of a short leg cast and a knee immobilizer. When properly applied, the knee immobilizer will combine effectively with the cast to immobilize both the knee and ankle. This technique can be used for a variety of knee fractures. Some fractures, however, require surgical repair and should have urgent orthopedic consultation.

Injury of the Supporting Structures of the Knee

Sprains involve injury to the ligaments of the knee. A Grade 1 sprain corresponds with a small incomplete tear and involves local tenderness with minimal swelling, no instability of the knee, and little increased pain with stress applied to the knee. A Grade 2 sprain corresponds with a moderate incomplete tear and involves local tenderness with moderate swelling, and some joint instability with a firm end-point on stress. A Grade 3 sprain cor-

Table 1. Ottawa Knee Rules

Plain radiographs are indicated if the patient has any one of the following conditions:

- Age 55 years or older
- Tenderness at head of fibula
- Isolated tenderness of the patella (no bone tenderness of knee other than patella)
- Inability to flex to 90°
- Inability to bear weight for four steps both in the ED and immediately after injury

Adapted from: Bachman LM, et al. The accuracy of the Ottawa knee rule to rule out knee fractures: A systematic review. *Ann Intern Med* 2004;140:121-124.

responds with a complete ligamentous disruption and involves local tenderness, swelling that can be minimal or marked, and significant joint instability with no clear end point.²⁹ (See Figure 9.)

Partial ligament tears typically are more painful and can produce significantly more swelling than complete disruptions. The MCL is the most commonly injured knee ligament. (See Figure 9.) The mechanism usually involves a strong lateral force applied to a slightly flexed knee. This type of injury also can produce ACL disruption and tearing of the medial meniscus. Isolated Grade 1 or Grade 2 sprains can be treated with immobilization, compression, rest, ice, elevation, and non-steroidal anti-inflammatory medications with orthopedic or primary care follow-up within a few days. Grade 3 injuries can be treated similarly with prompt orthopedic referral (within 24 hours).

The medial and lateral menisci also are commonly injured structures within the knee. They are not innervated, but injury may produce pain because of irritation from injury to adjacent ligaments. These structures are relatively avascular with their blood supply coming from peripheral capillaries. This poor blood supply results in slower accumulation of joint effusions associated with meniscal injury. The medial meniscus is fixed more securely and is more commonly injured. Acute meniscal tears are usually longitudinal and located peripherally.¹⁴ If it extends along the entire length of the meniscus, it is considered a bucket-handle tear. Traumatic ruptures are rare before age 10 years. They typically produce a triad of symptoms including joint line pain, swelling, and locking of the knee. The patient also may complain of the knee “giving out.” Treatment is the same as for a grade 1 or 2 sprain.

Knee trauma also can produce disruption or injury to the many tendinous insertions about the knee. Disruption of the quadriceps mechanism can occur in a number of possible locations. Such injuries commonly occur as the result of forced flexion of the knee with the quadriceps muscle contracted.³⁰ Rupture of the quadriceps tendon superior to the patella is more frequent in the elderly population, while patellar tendon ruptures inferior to the patella are more common in young athletes.³⁰ The extensor tendon mechanism also can be disrupted with a complete trans-

verse fracture of the patella and with avulsion of the tibial tubercle. Split and referral for orthopedic repair of the injury is recommended. Repeated, low-grade stress to the patella tendon may produce tendonitis or jumper’s knee.^{31,32}

Radiography

Plain radiography can be very helpful in the diagnosis of fracture, dislocation, and effusion about the knee after trauma; however, it reveals little useful information about ligamentous, meniscal, or tendon injuries.^{9,33-35} Standard views include a true anterior-posterior and lateral view of the knee along with one or two oblique views of the knee. Additional views may include a sunrise view of the knee, which is specifically used to identify patellar injuries.¹ In this projection, the knee is flexed, and the beam is directed tangentially to the longitudinal axis of the patella.

Historically, x-rays of the knee have been overused, and several clinical decision rules have been proposed to assist the examiner in determining which patients do not require imaging. The most widely accepted are the Ottawa Knee Rules,³⁶ which state that plain films of the knee are indicated if the patient is: 1) 55 years of age or older; 2) has tenderness at the head of the fibula; 3) has isolated tenderness of the patella; 4) is unable to flex the knee to 90°; or 5) is unable to bear weight for four steps both in the ED and immediately after the injury. (See Table 1.) These rules have helped decrease radiography rates, decrease cost to the patient, and decrease the length of stay in the ED without decreasing the rate of identifying clinically significant fractures of the knee.^{15,36}

Computerized tomography (CT) scanning of the knee occasionally is used by the orthopedic consultants to delineate further the anatomy of a fracture, but it has a limited role in the ED evaluation of the acutely injured knee.³⁷ Magnetic resonance imaging (MRI) is useful in the outpatient setting for evaluating the soft-tissue structures about the knee when the specific injury is in question, but usually is not necessary as part of the emergency assessment.^{9,33} When the type of injury is readily apparent from history and physical examination, the patient often will undergo arthroscopy without having an MRI.

Other Diagnostic Studies

Saline Arthrogram. When concerned about the possibility of a penetrating joint injury (e.g., a laceration that overlies the joint space), a saline arthrogram can be used safely to exclude communication with the joint. Proper performance of a saline arthrogram can help identify otherwise occult, open joint injuries.³⁸

To perform a saline arthrogram, the knee is prepped and draped in a sterile fashion, with the knee in a 30° flexion. Lidocaine 1% with or without epinephrine can be used as local anesthesia. A superior medial or lateral approach should be used similar to an arthrocentesis. The choice of approach is based upon avoiding the suspected area of penetration. Using the margin of the patella as a landmark, the clinician inserts an 18-gauge needle attached to a syringe containing sterile saline into the knee joint under negative pressure. Once placement is confirmed by aspiration of synovial fluid, 20-30 mL of saline is instilled into

the joint space, while the laceration or other skin injury is observed. Leakage of saline from the laceration confirms that there is communication with the joint space. Open joint injuries require emergent orthopedic consultation, and usually operative debridement and irrigation.

After finishing the procedure, be sure to remove all injected saline from the joint; the extra fluid may be quite uncomfortable for the patient.

Angiography. Angiography is a critical imaging modality used with suspected anterior dislocations, posterior dislocations, or bicruciate ruptures. Popliteal artery injury occurs in association with these injuries in 30-40% of cases.^{3,28} The prognosis is good when arterial injury is identified and repaired within 8 hours of injury. When the diagnosis is delayed, up to 86% of patients with popliteal artery injury require amputation, and 66% of those remaining have permanent ischemic changes of the leg and foot.^{16,21} If distal pulses are absent with this type of injury post-reduction, immediate surgical exploration is necessary.

Pitfalls

Several potential pitfalls exist in the evaluation of patients with knee injuries, including: 1) not getting a complete history regarding the mechanism of injury; 2) failure to have the knee re-examined in 24-48 hours when pain and swelling have decreased; 3) failure to examine the knee in flexion and extension; and 4) failure to consider popliteal artery injury in the face of significant knee injury, especially in the case of dislocations, and address this condition with either angiography or immediate surgical exploration.

Summary

Patients frequently present in the ED with trauma to the knee. It is critical for the ED physician to be familiar with the anatomy of the knee, trauma associated with certain mechanisms of injury, and appropriate diagnostic testing that is indicated based on the history and a thorough physical examination. Limitations of the initial examination should be recognized, as well as the more subtle findings of potential vascular compromise. Appropriate diagnostic testing should be performed and orthopedic consultation and follow-up obtained when indicated.

References

- Marx J, Hockberger R, Walls R, eds. *Rosen's Emergency Medicine: Concepts and Clinical Practice*. New York: Mosby; 2002:674-705.
- Tintinalli JE, Kelen GD, Stapczynski JS, et al, eds. *Emergency Medicine: A Comprehensive Study Guide*. New York: McGraw Hill; 2000:1814-1822.
- Seybold EA, Busconi BD. Traumatic popliteal artery thrombosis and compartment syndrome of the leg following blunt trauma to the knee: A discussion of treatment and complications. *J Orthop Trauma* 1996;10:138-141.
- Fadale PD, Noerdlinger MA. Sports injuries of the knee. *Curr Opin Rheumatol* 1999;11:144-150.
- Schenck RC Jr, Hunter RE, Ostrum RF, et al. Knee dislocations. *Instr Course Lect* 1999;48:515-522.
- Harner CD, Hoher J. Evaluation and treatment of posterior cruciate ligament injuries. *Am J Sports Med* 1998;26:471-482.
- Duri ZA, Aichroth P M, Zorrilla P. The posterior cruciate ligament: A review. *Am J Knee Surg* 1997;10:149-165.
- Mallory N, Kelsberg G, Ketchell D. Does a knee brace decrease recurrent ACL injuries? *J Fam Pract* 2003;52: 803-804.
- Ho CP, Marks PH, Steadman JR. MR imaging of knee anterior cruciate ligament and associated injuries in skiers. *Magn Reson Imaging Clin N Am* 1999;7:117-130.
- Smith BW, Green GA. Acute knee injuries: Part II. Diagnosis and management. *Am Fam Physician* 1995;51:799-806.
- Oeppen RS, Connolly SA, Bencardino JT, et al. Acute injury of the articular cartilage and subchondral bone: A common but unrecognized lesion in the immature knee. *AJR Am J Roentgenol* 2004; 182:111-117.
- Fritz RC. MR imaging of meniscal and cruciate ligament injuries. *Magn Reson Imaging Clin N Am* 2003;11: 283-293.
- Miller MD, Bergfeld JA, Fowler PJ, et al. The posterior cruciate ligament injured knee: Principles of evaluation and treatment. *Instr Course Lect* 1999;48:199-207.
- Swenson TM, Harner CD. Knee ligament and meniscal injuries. Current concepts. *Orthop Clin North Am* 1995;26:529-546.
- Shearman CM, el-Khoury GY. Pitfalls in the radiologic evaluation of extremity trauma: Part II. The lower extremity. *Am Fam Physician* 1998;57:1314-1322.
- Solomon DR, Simel DL, Bates DW, et al. Does this patient have a torn meniscus or ligament of the knee. *JAMA* 2003;286:1610-1620.
- Honari DB, Young CC, Boynton MD, et al. Diagnosis and treatment of common knee injuries in athletes. *Wis Med J* 1996;95:361-366.
- Muellner T, Nikolic A, Vecsei V. Recommendations for the diagnosis of traumatic meniscal injuries in athletes. *Sports Med* 1999;27: 337-345.
- Anderson AF, Lipscomb AB. Clinical diagnosis of meniscal tears. Description of a new manipulative test. *Am J Sports Med* 1986; 14:291-293.
- Noble J, Erat K. In defense of the meniscus: A prospective study of 200 meniscectomy patients. *J Bone Joint Surg Br* 1980;62:7-11.
- Zionts LE. Fractures around the knee in children. *J Am Acad Orthop Surg* 2002;10:345-355.
- Stover M. Distal femoral fractures: Current treatment, results and problems. *Injury* 2001;32: SC3-13.
- Iobst CA, Stanitski CL. Acute knee injuries. *Clin Sports Med* 2000;19:621-635.
- Dirschl DR, Dahners LE. Current treatment of tibial plateau fractures. *J South Orthop Assoc* 1997;6:54-61.
- Ruth JT. Fractures of the tibial plateau. *Am J Knee Surg* 2001;14: 125-128.
- Garner JP, Pike JM, George CD. Intra-articular dislocation of the patella: Two cases and literature review. *J Trauma* 1999;47:780-783.
- Ghalambor N, Vangsness CT Jr. Traumatic dislocation of the knee: A review of the literature. *Bull Hosp Jt Dis* 1995;54:19-24.
- Tominaga GT, Connolly JE, Wilson SE. Bilateral popliteal artery injury from bumper crush injury. *J Trauma* 1996;40:311-313.
- Perryman JR, Hershman EB. The acute management of soft tissue injuries of the knee. *Orthop Clin North Am* 2002;33:575-585.

30. Ilan DI, Tejwani N, Keschner M, et al. Quadriceps tendon rupture. *J Am Acad Orthop Surg* 2003;11:192-200.
31. Warden SJ, Brukner P. Patellar tendinopathy. *Clin Sports Med* 2003; 22:743-759.
32. Fredberg U, Bolvig L. Jumper's knee. Review of the literature. *Scand J Med Sci Sports* 1999;9:66-73.
33. Laughlin WJ Jr. Clinically appropriate, cost-effective MR scan of the acute knee injury. *J La State Med Soc* 1998;150: 308-311.
34. Tandeter HB, Shvartzman P. Acute knee injuries: Use of decision rules for selective radiograph ordering. *Am Fam Physician* 1999; 60:2599-2608.
35. Gray SD, Kaplan PA, Dussault R G. Imaging of the knee. Current status. *Orthop Clin North Am* 1997;28:643-658.
36. Bachmann LM, Haberzeth S, Steurer J, et al. The accuracy of the Ottawa knee rule to rule out knee fractures: A systematic review. *Ann Intern Med* 2004;140:121-124.
37. Bharam S, Vrahas MS, Fu FH. Knee fractures in the athlete. *Orthop Clin North Am* 2002;33:565-574.
38. Coumas JM, Palmer WE. Knee arthrography. Evolution and current status. *Radiol Clin North Am* 1998;36:703-728.

CE/CME Questions

1. Which of the following statements regarding the popliteal fossa is *not* true?
 - A. The popliteal fossa is a depression found behind the knee joint.
 - B. The popliteal artery represents the continuation of the femoral artery into the knee joint and is located in this recess.
 - C. The artery is not fixed within the fossa, which protects it from injury when the knee is stressed.
 - D. The peroneal nerve is located within this recess.
2. Which of the following statements is true regarding meniscal injuries?
 - A. Meniscal injuries are detected using the McMurray's test, Apley compression test or medial-lateral grind test.
 - B. The McMurray test involves flexion of the knee beyond 90° with the foot externally rotated in the examiner's hand. The other hand palpates the medial joint line as the knee is extended gradually beyond 90°. Pain, a "popping" sensation, or crepitus indicates a medial meniscus injury.
 - C. The McMurray test alone has been shown to have a low sensitivity and specificity for detecting medial meniscus injuries.
3. Which of the following statements regarding tibial fractures is *not* true?
 - A. Tibial fractures typically occur in one of two ways: 1) a direct blow to the tibia (bumper injury); or 2) axial compression of the tibia, which occurs when the knee is forcibly extended in anticipation of a frontal impact.
 - B. Class A injuries (condylar and plateau fractures) usually are produced by axial loading (e.g., a fall from a height).
 - C. The rotational stress on the knee will determine whether the medial or lateral tibial plateau is affected.
 - D. The medial plateau is the site of fracture in the majority of cases.
4. Which of the following statements regarding disruption of the quadriceps mechanism is true?
 - A. Such injuries commonly occur as the result of forced extension of the knee with the quadriceps muscle contracted.
 - B. Rupture of the quadriceps tendon superior to the patella is more frequent in the elderly population, while patellar tendon ruptures inferior to the patella are more common in young athletes.
 - C. The abduction tendon mechanism also can be disrupted with a complete transverse fracture of the patella and with avulsion of the tibial tubercle.
 - D. Immediate arteriography is indicated.
5. Based upon the Ottawa rules, in which of the following patients should plain radiographs *not* be obtained?
 - A. A 65-year-old male who has fallen down the steps and complains of knee pain.
 - B. A 42-year-old female who presents after falling at a football game with tenderness at the head of the fibula.
 - C. A 37-year-old skier who presents after falling and has isolated tenderness of the patella.
 - D. A 52-year-old male who presents with knee pain and can flex his knee to 90°.

CME Objectives

Upon completing this program, the participants will be able to:

- a.) Quickly recognize or increase index of suspicion for occult knee injury;
- b.) Be educated about how to correctly and quickly stabilize, and then to manage patients with knee trauma;
- c.) Understand various diagnostic modalities for arterial injuries; and
- d.) Understand both likely and rare complications that may occur.

CE/CME Instructions

Physicians and nurses participate in this continuing medical education/continuing education program by reading the article, using the provided references for further research, and studying the questions at the end of the article. Participants should select what they believe to be the correct answers, then refer to the list of correct answers to test their knowledge. To clarify confusion surrounding any questions answered incorrectly, please consult the source material. **After completing this activity, you must complete the evaluation form provided and return it in the reply envelope provided in order to receive a certificate of completion.** When your evaluation is received, a certificate will be mailed to you.

6. Which of the following statements regarding knee dislocations is *not* true?
 - A. Knee dislocation is the result of a significant force delivered to the knee and occurs as the result of hyperextension, posterior forces to the tibia, or significant rotary forces.
 - B. Any knee dislocation is associated with significant ligamentous disruption.
 - C. There are three types of dislocation of the knee: anterior, posterior, and lateral.
 - D. In anterior dislocations (the most common type), the tibia is displaced anteriorly to the femur and typically is the result of hyperextension of the knee joint.
 - E. In a lateral dislocation, the tibia is forced posteriorly by a direct force applied to the proximal tibia.
7. Which of the following statements regarding lateral dislocations is *not* true?
 - A. Lateral dislocations can be medial or lateral and occur with equal frequency
 - B. Lateral dislocations can be complicated further with rotary deformity of the knee.
 - C. Any type of knee dislocation carries a 75% incidence of open joint injury.
 - D. Reduction of the dislocation requires longitudinal traction.
 - E. Sedation and pain relief commonly are required.
8. Which of the following statements regarding vascular injuries is true?
 - A. Injury to the popliteal artery is common in knee dislocations (up to 40% of all dislocations).
 - B. It is critical that such injuries be identified early and repaired within 6 to 8 hours of injury.
 - C. Intact pulses in the foot do not exclude arterial injury.

- D. An arteriogram should be performed on these patients according to most experts, although an alternative strategy of admission and repeated clinical examination after normal ankle brachial indices (ABI) in the ED has been suggested.
 - E. All of the above statements are true.
9. Which of the following statements regarding an ABI is *not* true?
 - A. An ABI is performed by obtaining a blood pressure in the arm by standard technique, then taking a blood pressure at the ankle. Dividing the ankle pressure by the arm pressure determines the ABI.
 - B. Ankle pressure should be greater than brachial pressure.
 - C. A ratio of one or greater is considered normal.
 - D. An ABI of 0.9 or less always means that the patient has sustained an arterial injury.
 - E. Patients who have peripheral arterial disease also may have an ABI of less than 0.9.
10. A 35-year-old male presents with a patellar fracture. Which of the following statements is true?
 - A. A patellar fractures typically is caused by direct knee trauma.
 - B. A secondary mechanism may involve an avulsion caused by the pull of the quadriceps mechanism.
 - C. Patellar fractures are more common in males.
 - D. A non-displaced fracture can be treated with a knee immobilizer and pain control.
 - E. All of the above statements are true.


Answer Key:

1. C; 2. B; 3. D; 4. B; 5. D; 6. E; 7. C; 8. E; 9. D; 10. E

In Future Issues:

Ocular Trauma

A new resource perfect for heat-of-battle patient care



**ORDER YOUR COPY
FOR ONLY \$79
(BOOK # 504115)**

You face hundreds of trauma patients every week; finding quick and accurate answers to your critical questions shouldn't be another traumatic experience. That's why *Thomson American Health Consultants* offers you this new pocket-sized guide.

"Trauma Reports: Rapid Reference Guidelines" gets right to the point. Your job is too hectic for anything else.

You get at-a-glance treatment strategies in a summarized format. Our most popular, and heavily-used "guideline cards" are all pulled together into this single pocket guide. You get proven effective...

- diagnosis and treatment charts
- algorithms
- treatment guidelines

Because the topics covered in this pocket guide cover the situations you encounter every day, the facts are immediately applicable! You get the added confidence of always having relevant, clinically accurate answers in the pocket of your lab coat.

**Call (800) 688-2421 (404 262-5476 outside the US and Canada) or visit our website
at www.ahcpub.com. Please refer to code 61182.**